

Addendum to “On the p^λ problem”

Stephan Baier *

February 2, 2008

Abstract: We prove that the conditions $\lambda < 5/19$ and $L \leq T^{1/2}$ in Theorems 3 and 4 of our recent paper [Bai] can be omitted.

In [Bai] we proved the following mean value estimate for products of shifted and ordinary Dirichlet polynomials.

THEOREM 1: ([Bai, Theorem 4]) *Suppose that $\alpha \neq 0$, $0 \leq \theta < 1$, $T > 0$, $K \geq 1$, $L \geq 1$. If $\theta \neq 0$, then additionally suppose that $L \leq T^{1/2}$. Let (a_k) and (b_l) be arbitrary sequences of complex numbers. Suppose that $|a_k| \leq A$ for all $k \sim K$ and $|b_l| \leq B$ for all $l \sim L$. Then,*

$$(1) \quad \int_0^T \left| \sum_{k \sim K} a_k k^{it} \right|^2 \left| \sum_{l \sim L} b_l (l + \theta)^{i\alpha t} \right|^2 dt \\ \ll A^2 B^2 (T + KL) KL \log^3(2KLT),$$

the implied $\ll$ -constant depending only on α . If $\theta = 0$, then $\log^3(2KLT)$ on the right side of (1) may be replaced by $\log^2(2KLT)$.

We then used this mean value estimate to prove the following result on the p^λ problem.

THEOREM 2: ([Bai, Theorem 3]) *Suppose that $\varepsilon > 0$, $B > 0$, $\lambda \in (0, 1/2]$ and a real θ are given. If θ is irrational, then suppose that $\lambda < 5/19$.*

*e-mail: sbaier@mast.queensu.ca

Let $N \geq 3$. Let $\mathbf{A}$ be an arbitrarily given subset of the set of positive integers. Define

$$F_\theta(\lambda) := \begin{cases} \max_{k \in \mathbf{N}} \min \left\{ \frac{5}{12} - \frac{(k+6)\lambda}{6(k+1)}, \frac{5}{11} - \frac{(5k+1)\lambda}{11} \right\} & \text{if } \theta \text{ is rational,} \\ \frac{5}{12} - \frac{7\lambda}{12} & \text{otherwise.} \end{cases}$$

Suppose that

$$N^{-F_\theta(\lambda) + \varepsilon \lambda} \leq \delta \leq 1.$$

Then we have

$$\sum_{\substack{N < n \leq 2N, \\ \{n^\lambda - \theta\} < \delta, \\ [n^\lambda] \in \mathbf{A}}} \Lambda(n) = \frac{\delta}{\lambda} \cdot \sum_{\substack{N^\lambda < n \leq (2N)^\lambda, \\ n \in \mathbf{A}}} n^{1/\lambda - 1} + O\left(\frac{\delta N}{(\log N)^B}\right).$$

At the end of the last section in [Bai] we pointed out that if the condition $L \leq T^{1/2}$ in the above Theorem 1 could be omitted, then the condition $\lambda < 5/19$ in Theorem 2 could be omitted too. In the following we will see that the condition $L \leq T^{1/2}$ in Theorem 1 is actually superfluous if we allow ourselves to weaken the mean value estimate (1) slightly. We establish the following

THEOREM 3: Let $\theta, \xi, \alpha, \beta$ be real numbers with $0 \leq \theta, \xi < 1$ and $\alpha\beta \neq 0$. Suppose that $T, K, L \geq 1$, $|a_k| \leq 1$ and $|b_l| \leq 1$. Then

$$(2) \quad \int_0^T \left| \sum_{k \sim K} a_k (k + \theta)^{i\alpha t} \right|^2 \left| \sum_{l \sim L} b_l (l + \xi)^{i\beta t} \right|^2 dt \ll (T + KL) KL (\log T)^{15}.$$

In accordance with the proof of [Bai, Theorem 3], from the above Theorem 3 with $\xi = 0$ it can be deduced that Theorem 2 holds true with the condition $\lambda < 5/19$ omitted.

The main idea of our proof of Theorem 3 is to relate the shifted Dirichlet polynomials on the left-hand side of (2) to the corresponding Hurwitz zeta functions. For technical reasons we here define the Hurwitz zeta function $\zeta(s, y)$ in a slightly different manner to normal usage. For $0 \leq y < 1$ and $\text{Re } s > 1$ we write

$$\zeta(s, y) := \sum_{n=1}^{\infty} (n + y)^{-s}.$$

In the usual definition the series on the right-hand side starts with $n = 0$, and the case $y = 0$ is excluded, which we seek to avoid here.

As a function of s , the Hurwitz zeta function has a meromorphic continuation to the entire complex plane, with a simple pole at $s = 1$ (see [Ivi]). At first, we establish the following fourth power moment estimate for the Hurwitz zeta function on the critical line.

THEOREM 4: *Suppose that $V > 2\pi$ and $0 \leq y < 1$. Then*

$$\int_{-V}^V \left| \zeta \left(\frac{1}{2} + it, y \right) \right|^4 dt \ll V(\log V)^{10}.$$

PROOF: By $\zeta(\bar{s}, y) = \overline{\zeta(s, y)}$, it suffices to show that

$$(3) \quad \int_{2\pi}^V \left| \zeta \left(\frac{1}{2} + it, y \right) \right|^4 dt \ll V(\log V)^{10}.$$

By [Tch, Lemma 1], the Hurwitz zeta function satisfies an approximate functional equation of the form

$$\begin{aligned} \zeta \left(\frac{1}{2} + it, y \right) &= \sum_{1 \leq m \leq M} (m + y)^{-1/2 - it} + \chi(1/2 + it) \sum_{1 \leq n \leq N} e(-ny) n^{-1/2 + it} + \\ &O \left(1 + M^{-3/2} |t|^{1/2} \right) \end{aligned}$$

if $|t| \geq 2\pi$, $1 \leq M \leq |t|$, $N \geq 1$ and $2\pi MN = |t|$, where $|\chi(1/2 + it)| = 1$. Hence, we have

$$\begin{aligned} (4) \quad \int_{2\pi}^V \left| \zeta \left(\frac{1}{2} + it, y \right) \right|^4 dt &\ll V + \int_{2\pi}^V \left| \sum_{1 \leq m \leq \sqrt{t/(2\pi)}} (m + y)^{-1/2 - it} \right|^4 dt \\ &+ \int_{2\pi}^V \left| \sum_{1 \leq n \leq \sqrt{t/(2\pi)}} e(-ny) n^{-1/2 + it} \right|^4 dt. \end{aligned}$$

By the orthogonality relation

$$\int_0^1 e(zu) du = \begin{cases} 1 & \text{if } z = 0, \\ 0 & \text{if } z \in \mathbb{Z} \setminus \{0\}, \end{cases}$$

we get

$$(5) \quad \sum_{1 \leq m \leq \sqrt{t/(2\pi)}} (m+y)^{-1/2-it} = \int_0^1 \sum_{1 \leq m \leq \sqrt{V}} (m+y)^{-1/2-it} e(mu) K(t, u) \, du$$

for $2\pi \leq t \leq V$, where

$$K(t, u) := \sum_{1 \leq n \leq \sqrt{t/(2\pi)}} e(-nu).$$

If $2\pi \leq t \leq V$, then the geometric sum $K(t, u)$ can be estimated by

$$(6) \quad K(t, u) \ll \min\{\sqrt{V}, ||u||^{-1}\}.$$

This yields

$$(7) \quad \int_0^1 |K(t, u)| \, du \ll \log V.$$

Using Hölder's inequality, from (5) and (7), we obtain

$$(8) \quad \left| \sum_{1 \leq m \leq \sqrt{t/(2\pi)}} (m+y)^{-1/2-it} \right|^4 \ll (\log V)^3 \int_0^1 \left| \sum_{1 \leq m \leq \sqrt{V}} (m+y)^{-1/2-it} e(mu) \right|^4 |K(t, u)| \, du.$$

Employing Hölder's inequality and [Har, Lemma 3] after dividing the sum on the right-hand side of (8) into $O(\log V)$ sums of the form

$$\sum_{M < m \leq 2M} (m+y)^{-1/2-it} e(mu),$$

we obtain

$$(9) \quad \int_{2\pi}^V \left| \sum_{1 \leq m \leq \sqrt{V}} (m+y)^{-1/2-it} e(mu) \right|^4 dt \ll V(\log V)^6,$$

where the implied $\ll$ -constant does not depend on u . Combining (6), (8) and (9), we get

$$(10) \quad \int_{2\pi}^V \left| \sum_{1 \leq m \leq \sqrt{t/(2\pi)}} (m+y)^{-1/2-it} \right|^4 dt \ll V(\log V)^{10}.$$

In a similar manner, we can prove

$$(11) \quad \int_{2\pi}^V \left| \sum_{1 \leq n \leq \sqrt{t/(2\pi)}} e(-ny)n^{-1/2+it} \right|^4 dt \ll V(\log V)^{10}.$$

Combining (4), (10) and (11), we obtain (3). This completes the proof. $\square$

To all appearances, there is no result like Theorem 4 in the literature.

We now prove Theorem 3 along the lines of the proof of [BaH, Theorem 3]. First we write

$$F(t) := \sum_{k \sim K} a_k(k+\theta)^{it}, \quad G(t) := \sum_{l \sim L} b_l(l+\xi)^{it}, \quad D(t) := \sum_{k \sim K} (k+\theta)^{it},$$

$$E(t) := \sum_{l \sim L} (l+\xi)^{it}.$$

Similar as in the proof of [BaH, Theorem 3], we can suppose that $K \leq L \leq T$, for otherwise the desired estimate follows from a classical mean value estimate for $G(t)$.

Analogous to [BaH, (17)], we have

$$(12) \quad \int_0^T |F(\alpha t)G(\beta t)|^2 dt \ll (KL)^2 + \log T \max_{1 \leq V \leq T} \int_V^{2V} |D(\alpha t)E(\beta t)|^2 dt.$$

We fix V in the interval $1 \leq V \leq T$ for which the maximum is attained.

In the same manner like [BaH, (19)] one can prove

$$(13) \quad |D(\alpha t)| \ll K^{1/2} \int_{-V}^V \left| \zeta \left(\frac{1}{2} + i\sigma - i\alpha t, \theta \right) \right| \rho(\sigma) d\sigma + \frac{K \log T}{V}$$

as well as

$$(14) \quad |E(\beta t)| \ll L^{1/2} \int_{-V}^V \left| \zeta \left(\frac{1}{2} + i\tau - i\beta t, \xi \right) \right| \rho(\tau) d\tau + \frac{L \log T}{V},$$

where $\rho(x) := \min(1, 1/|x|)$. Using (13), (14) and the inequality of Cauchy-Schwarz, we deduce

$$(15) \quad \begin{aligned} & \int_V^{2V} |D(\alpha t) E(\beta t)|^2 dt \\ & \ll \frac{(KL)^2 \log^4 T}{V^3} + \frac{K^2 L \log^3 T}{V^2} \int_{-V}^V \rho(\tau) \int_V^{2V} \left| \zeta \left(\frac{1}{2} + i\tau - i\beta t, \xi \right) \right|^2 dt d\tau + \\ & \quad \frac{KL^2 \log^3 T}{V^2} \int_{-V}^V \rho(\sigma) \int_V^{2V} \left| \zeta \left(\frac{1}{2} + i\sigma - i\alpha t, \theta \right) \right|^2 dt d\sigma + KL \log^2 T \times \\ & \quad \int_{-V}^V \int_{-V}^V \rho(\sigma) \rho(\tau) \int_V^{2V} \left| \zeta \left(\frac{1}{2} + i\sigma - i\alpha t, \theta \right) \zeta \left(\frac{1}{2} + i\tau - i\beta t, \xi \right) \right|^2 dt d\sigma d\tau \\ & \ll \frac{(KL)^2 \log^4 T}{V^3} + \frac{K^2 L \log^4 T}{V^2} \int_{-CV}^{CV} \left| \zeta \left(\frac{1}{2} + it, \xi \right) \right|^2 dt + \\ & \quad \frac{KL^2 \log^4 T}{V^2} \int_{-CV}^{CV} \left| \zeta \left(\frac{1}{2} + it, \theta \right) \right|^2 dt + KL \log^4 T \times \\ & \quad \left(\int_{-CV}^{CV} \left| \zeta \left(\frac{1}{2} + it, \theta \right) \right|^4 dt \right)^{1/2} \left(\int_{-CV}^{CV} \left| \zeta \left(\frac{1}{2} + it, \xi \right) \right|^4 dt \right)^{1/2}, \end{aligned}$$

where C is a certain constant which depends only on α and β . From (12), (15), Theorem 4 and a similar *second* power moment estimate for the Hurwitz zeta function (which can be derived directly from Theorem 4 using the inequality of Cauchy-Schwarz), we obtain (2). This completes the proof of

Theorem 3. $\square$

Acknowledgements. This research has been supported by a Marie Curie Fellowship of the European Community programme “Improving the Human Research Potential and the Socio-Economic Knowledge Base” under contract number HPMF-CT-2002-02157.

References

- [Bai] S. Baier, *On the p^λ problem*, Acta Arith. 113 (2004), 77-101.
- [BaH] A. Balog, G. Harman, *On mean values of Dirichlet polynomials*, Arch. Math. (Basel) 57 (1991), 581-587.
- [Har] G. Harman, *Fractional and integral parts of p^λ* , Acta Arith. 58 (1991), 141-152.
- [Ivi] A. Ivic, *The Riemann Zeta-Function*, Wiley-Interscience, New York 1985.
- [Tch] N. Tchudakoff, *On Goldbach-Vinogradov’s theorem*, Ann. Math. 48 (1947), 515-545.